

# Maple Syrup Urine Disease (MSUD)

Biochemistry 5614 Service-Learning Initiative Autumn 2020

Instructor: Dr. Venkat Gopalan

Rosemary Zaleski, Gianni Giarrano, Andrew Sobodosh, Mia Kordowski,  
Halli Petillo, Owen Newlove, Zechariah Shearer, Josh Gilger, Claudia Studebaker




# Occurrence frequency

- United States population
  - Estimated frequency of approximately 1 in 220,000 live births
- The Mennonite population (United States)
  - Estimated frequency of approximately 1 in 380 live births

(The much higher frequency among the Mennonites is due to the founder effect, i.e., when inbreeding in isolated groups results in a small gene pool with persistence of genes/mutation)
- A child may have increased risk of MSUD due to both parents being carriers of a spontaneous/random mutation, of Mennonite lineage, or both

# Genetics


- Autosomal recessive disorder (i.e., two mutant copies needed to manifest disease)
- The mutation could be in one of three genes that code for the branched chain  $\alpha$ -keto-acid dehydrogenase complex (BCKD), which breaks down three branched chain amino acids



Digital image retrieved Apr 7, 2021

from <https://www.mayoclinic.org/tests-procedures/genetic-testing/multimedia/genetic-disorders/sls-20076216?s=4>

# Genetics overview video



<https://youtu.be/Nv6qUsKYodA>



# Major food groups\*


## Carbohydrates



## Fats



## Proteins



- \* Each exemplar shown above was chosen because of its representative value to the respective group but note that the depicted item is not entirely devoid of the two other food groups

**MSUD associated with a  
defect in protein  
utilization**

# Nature of the problem in MSUD


Amino acids are the building blocks of proteins

In MSUD, defective BCKD enzyme cannot break down these three essential\* amino acids




## BRANCHED CHAIN AMINO ACIDS


*Leucine*



*Isoleucine*



*Valine*






Digital image retrieved Nov 7, 2020 from  
<https://killcliff.com/blogs/hpc/do-you-need-bcaas>

\* Essential refers to those amino acids that the body cannot make and must therefore be obtained from the diet




# Biochemical features

- Defective BCKD enzyme – variants of MSUD have varying severity of deficits in activity
- Build-up of branched chain amino acids and  $\alpha$ -keto acids (reflected in high levels in blood and urine)
- Stools, urine, sweat, and earwax that smells of maple syrup, the odor associated with  $\alpha$ -keto acids

## Branched Chain Amino Acids (BCAAs)



## $\alpha$ -keto acids



# Symptoms


- Symptoms can appear starting about 3 days after birth as the toxic metabolites accumulate; symptoms include:
  - **Bodily fluids smell of maple syrup**
  - Poor feeding, vomiting, loss of appetite, irritability
  - Tiredness and weakness
  - Poor muscle tone ("floppy") or stiffness of muscles
  - Abnormal muscle movements
  - Developmental delay
  - Seizures and convulsions leading to respiratory failure/coma
- Severity of symptoms may be dependent upon which clinical variant of MSUD an individual has





# Diagnosis

- Birth
  - Newborn screening
  - May result in a false negative due to lack of adequate time for measurable accumulation of toxic metabolites
- Post birth/toddler
  - Analysis of plasma, urine, and white blood cells, symptom assessment
  - Amino acid and BCKD enzyme testing to check for elevated levels




## Newborn screening

Digital image retrieved Nov 07, 2020 from  
<https://www.af.mil/News/Photos/igphoto/2000419731/>

# Prenatal diagnosis

If a fetus is predisposed to MSUD (based on either family history or previously affected child), two prenatal diagnosis methods can be used for testing of BCKD enzyme activity:

- **Chorionic villus sampling:** performed at 11-14 weeks; biopsy of cells from mother's placenta for testing; low risk of amniotic leakage or of infection
- **Amniocentesis:** performed at 15-20 weeks; amniotic fluid is removed and tested; some risk of amniotic leakage and infection




## Amniocentesis

Digital image retrieved Nov 7, 2020 from  
<https://www.mayoclinic.org/tests-procedures/amniocentesis/about/pac-20392914>

# Therapy

- Strict high-calorie diet to prevent breakdown of body's energy stores and weight loss
- Limit intake of leucine, isoleucine, and valine (present in all proteins); intake is limited for infants by specialized formulas
  - Underestimating intake may result in development of symptoms
- Frequently monitor levels of BCAAs in the blood (twice a week for infants and weekly for older children/adults)
- Liver transplant to restore functional enzyme; albeit a possibility, this approach has its own risks, and it will not prevent the defective MSUD gene to be passed onto children



\* An individual with MSUD must limit dietary intake of these food types to avoid a metabolic crisis

# Prognosis

- Maple Syrup Urine Disease is managed through strict diet
  - BCAA-free formula for infants provides necessary dietary sustenance without harmful BCAAs
  - BCAAs are supplemented into the diet in very low amounts
  - However, even with strict diet control, an individual is still at high risk for metabolic attack
- An affected individual can grow into adulthood with normal development
- Symptoms will vary in intensity based on the relative activity of the enzyme in affected individuals



## MSUD Formula

Digital image retrieved Nov 07, 2020 from  
<https://www.msudsupport.org/newsletters/116-volume-33-2/619-introducing-new-infant-formula-for-msud>



# Available support

## [MSUD Family Support Group](#)

- Access to current, relevant MSUD research
- Network of MSUD-specific dietitians, physicians, and facilities




## [Organic Acidemia Association](#)

- Information on diagnosis procedures
- Facebook support group, conferences, and research fundraising

## [National Organization for Rare Disorders](#)

- Patient and family advocacy
- Information for financial assistance and research opportunities



# References

- Allen, R. J. (2009, July 17). A Brief History of Maple Syrup Urine Disease (MSUD). Retrieved Nov. 7, 2020, from <https://www.msud-support.org/newsletters/13-volume-11-2/41-a-brief-history-of-maple-syrup-urine-disease-msud>
- Amniocentesis - Mayo Clinic*. (2019, March 8). Mayo Clinic. Digital Image Retrieved Nov. 7, 2020 from <https://www.mayoclinic.org/tests-procedures/amniocentesis/about/pac-20392914>
- Blackburn, P. R., Gass, J. M., Vairo, F., Farnham, K. M., Atwal, H. K., Macklin, S., Klee, E. W., & Atwal, P. S. (2017). Maple syrup urine disease: mechanisms and management. *The application of clinical genetics*, 10, 57–66. <https://doi.org/10.2147/TACG.S125962>
- Buist, N. R., & Zupan, M. (2020, June 05). Maple Syrup Urine Disease. Retrieved October 21, 2020, from <https://rarediseases.org/rare-diseases/maple-syrup-urine-disease/>
- Chapman, K. A., Gramer, G., Viall, S., & Summar, M. L. (2018). Incidence of maple syrup urine disease, propionic acidemia, and methylmalonic aciduria from newborn screening data. *Molecular genetics and metabolism reports*, 15, 106–109. <https://doi.org/10.1016/j.ymgmr.2018.03.011>
- Chorionic Villus Sampling. (2020, Nov 12). Mayo Clinic. Retrieved April 8, 2021 from <https://www.mayoclinic.org/tests-procedures/chorionic-villus-sampling/about/pac-20393533>
- Cliff, K. (2020, April 03). *Do you need BCAAs?* Kill Cliff. <https://killcliff.com/blogs/hpc/do-you-need-bcaas>
- Home - MSUD Family Support Group. (n.d.). Retrieved Nov. 7, 2020, from <https://www.msud-support.org/>
- Home - NORD. (n.d.). Retrieved Nov. 7, 2020, from <https://rarediseases.org/>
- Home – Organic Acidemia Association. (n.d.). Retrieved Nov. 10, 2020, from <https://www.oaanews.org/>
- Maple Syrup Urine Disease. (2020, June 05). NORD (National Organization for Rare Disorders). <https://rarediseases.org/rare-diseases/maple-syrup-urine-disease/>
- Maple syrup urine disease: Management and treatment. (18 Nov 2019). *Cleveland Clinic*. Retrieved on October 28, 2020 from <https://my.clevelandclinic.org/health/diseases/21168-maple-syrup-urine-disease/management-and-treatment>
- Strauss KA, Puffenberger EG, Carson VJ. Maple Syrup Urine Disease. 2006 Jan 30 [Updated 2020 Apr 23]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2020. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK1319/>
- Wappner, R. S. (2009, July 20). Description and treatment of maple syrup urine disease. *MSUD Family Support Group*. Retrieved on October 28, 2020 from <https://www.msud-support.org/newsletters/33-volume-19-1/219-description-a-treatment-of-maple-syrup-urine-disease>

# Contributors

Name	Project Role
Rosemary Zaleski	Available support and text lead
Gianni Giarrano	Prognosis and figures lead
Andrew Sobodosh	Occurrence frequency
Mia Kordowski	History and provenance
Halli Petillo	Therapy
Owen Newlove	Biochemical features
Zechariah Shearer	Genetics
Josh Gilger	Symptoms
Claudia Studebaker	Diagnosis